Author Index

Ahsan, T.
——, Colenutt, B. A. and Sing, K. S. W.
Gas chromatography of pure and surface-modified precipitated calcium carbonate 17
Albert, K., see Hetem, M. 269
Albrecht, W., see Engel, K.-H. 176
Artz, W. E., see Schmitz, H. H. 261
Ashraf, M., see Singh, A. K. 233
Avilés, F. X., see Serra, M. A. 27
Bardarov, V.
——, Zaikov, Chr. and Mitewa, M.
Application of high-performance liquid chromatography with spectrophotometric and electrochemical detection to the analysis of alkylenebis(dithiocarbamates) and their metabolites 97
Bayer, E., see Hetem, M. 269
Brand, F., see Simon, P. 445
Brorson, T., see Skarping, G. 125
Brown, F. R.
—— and Draper, W. M.
Separation of phenols and their glucuronide and sulfate conjugates by anion-exchange liquid chromatography 441
Buchs, A., see Gülaçar, F. O. 61
Burke, III, J. A., see Pirkle, W. H. 377
Cardwell, T. J.
—— and Lorman, T. H.
Liquid chromatographic analysis of bromination reactions of metal trifluoroacetylacetonates 181
Chang, J.-P., see Pirkle, W. H. 377
Chikazumi, N.
——, Mukoyama, Y. and Sugitani, H.
High-performance size-exclusion chromatography of poly- and oligoethylene terephthalate using a mixture of hexafluoroisopropanol and chloroform as the mobile phase 85
Colenutt, B. A., see Ahsan, T. 17
Cooper, R. S., see Warth, L. M. 401
Copsey, J., see Houghton, E. 73
Cramers, C., see Hetem, M. 269
Cuchillo, C. M., see Serra, M. A. 27
Cuhra, P., see Hajšlová, J. 243
Dalene, M., see Sandström, J. F. 135
——, see Skarping, G. 125
——, see Tiljander, A. 145
Davídek, J., see Hajšlová, J. 243
Davídek, T., see Hajšlová, J. 243
De Haan, J., see Hetem, M. 269
Dhara, K. P., see Lewis, N. G. 345
Dietz, J. M., see Schmitz, H. H. 261
Dios, C. G., see Sánchez-Rasero, F. 424
Dombrovskis, D., see Singh, A. K. 233
Draper, W. M., see Brown, F. R. 441
Dubin, P. L.
—— and Principi, J. M.
Optimization of size-exclusion separation of proteins on a Superose column 159
Dumasia, M. C., see Houghton, E. 73
Eckert, H., see Müller, H. 452
Emery, M. F.
—— and Lim, C. K.
Separation of cationic technetium-99m amine complexes on porous graphitic carbon 212
Engel, K.-H.
——, Flath, R. A., Albrecht, W. and Tressl, R.
Gas chromatographic separation of diastereomeric dicarbamate derivatives of γ- and δ-lactones 176
Erdman, Jr., J. W., see Schmitz, H. H. 261
Evans, M. B.
——, Smith, M. S. and Oxford, J. M.
Use of a stop-flow technique to study on-column decomposition in supercritical fluid chromatography 170
Ezawa, T., see Nakai, S. 459
Falk, O., see Jacobsson, S. 194
Fielden, P. R.
—— and Packham, A. J.
Selective determination of benzo[*a*]pyrene in petroleum-based products using multi-column liquid chromatography 117
Flath, R. A., see Engel, K.-H. 176
Fritz, J. S., see Warth, L. M. 401
Fujii, Y.
——, Ikeda, Y. and Yamazaki, M.
High-performance liquid chromatographic determination of secondary cardiac glycosides in *Digitalis purpurea* leaves 319
Gandhe, B. R.
——, Malhotra, R. C. and Gutch, P. K.
Gas chromatographic retention indices of tear gases on capillary columns 165
Ginn, A., see Houghton, E. 73
Giralt, E., see Serra, M. A. 27
Gordon, B., see Singh, A. K. 233
Granley, K., see Singh, A. K. 233
Gülaçar, F. O.
——, Buchs, A. and Susini, A.
Capillary gas chromatography–mass spectrometry and identification of substituted carboxylic acids in lipids extracted from a 4000-year-old Nubian burial 61
Gutch, P. K., see Gandhe, B. R. 165
Haan, J. de, see Hetem, M. 269

Hajšlová, J.
——, Cuhra, P., Davídek, T. and Davídek, J.
Gas chromatographic determination of diquat and paraquat in crops 243
Hällgren, J.-E., see Wingsle, G. 335
Hermans-Lokkerbol, A.
——, Van der Leer, T. and Verpoorte, R.
Reversed-phase high-performance liquid chromatographic separation of some indole and quinoline alkaloids from *Cinchona* 39
Hetem, M.
——, Van de Ven, L., De Haan, J., Cramers, C., Albert, K. and Bayer, E.
Study of the changes in mono-, di- and trifunctional octadecyl-modified packings for reversed-phase high-performance liquid chromatography with different eluent compositions 269
Hewetson, D., see Singh, A. K. 233
Hill, Jr., H. H., see St. Louis, R. H. 221
Houghton, E.
——, Ginn, A., Teale, P., Dumasia, M. C. and Copsey, J.
Comparison of the use of mass spectrometry and methylene unit values in the determination of the stereochemistry of estranediol, the major urinary metabolite of 19-nortestosterone in the horse 73
Hsu, F.-F.
—— and Sherman, W. R.
Performing high salt concentration gradient elution ion-exchange separations using thermospray mass spectrometry 437
Hurtubise, R. J., see Olsen, L. D. 5
Ikeda, Y., see Fujii, Y. 319
Inciong, Ma. E. J., see Lewis, N. G. 345
Ito, Y., see Oka, H. 53
Jacobsson, S.
—— and Falk, O.
Determination of hydrogen sulphide by porous-layer open-tubular column gas chromatography–mass spectrometry 194
Janda, V.
——, Steenbeke, G. and Sandra, P.
Supercritical fluid extraction of *s*-triazine herbicides from sediment 200
Jenke, D. R.
Effect of some operational variables on the efficiency of ion chromatographic separations 387
Kerby, N. W.
——, Reed, R. H. and Rowell, P.
Separation of algal organic osmolytes by high-performance liquid chromatography 353
Kleinmann, I.
——, Plicka, J., Šmídl, P. and Svoboda, V.
Hydrophobic interaction chromatography of proteins on Separon HEMA. I. The effect of an initial salt concentration on the separation of proteins 327
Kobayashi, T., see Nakai, S. 459
Kokotti, H., see Pukkila, J. 369
Krob, E. J., see Walter, H. 307
Leer, T. van der, see Hermans-Lokkerbol, A. 39
Lemacon, C., see Simon, P. 445
Leung, C. P.
—— and Leung, W. K. H.
Determination of vitamin U and its degradation products by high-performance liquid chromatography with fluorescence detection 361
Leung, W. K. H., see Leung, C. P. 361
Lewis, N. G.
——, Inciong, Ma. E. J., Dhara, K. P. and Yamamoto, E.
High-performance liquid chromatographic separation of *E*- and *Z*-monolignols and their glucosides 345
Lim, C. K., see Emery, M. F. 212
Lorman, T. H., see Cardwell, T. J. 181
McCune, J. E., see Pirkle, W. H. 419
Malhotra, R. C., see Gandhe, B. R. 165
Mesbah, M.
—— and Whitman, W. B.
Measurement of deoxyguanosine/thymidine ratios in complex mixtures by high-performance liquid chromatography for determination of the mole percentage guanine + cytosine of DNA 297
Metz, P. A.
——, Morse, F. L. and Theyson, T. W.
Quantification and characterization of the trifluoroacetic anhydride derivatives of N,N′-ethylenebisstearamide and N,N′-ethylenebisoleamide 107
Mischke, C. F., see Norman, H. A. 206
Mishra, U., see Singh, A. K. 233
Mitewa, M., see Bardarov, V. 97
Morse, F. L., see Metz, P. A. 107
Mukoyama, Y., see Chikazumi, N. 85
Müller, H.
—— and Eckert, H.
Simultaneous determination of monoethanolamine and glycine betaine in plants 452
Nakai, S.
——, Kobayashi, T. and Ezawa, T.
Photodecomposition of Moclobemide on a silica gel thin-layer chromatographic plate 459

Norman, H. A.
——, Mischke, C. F. and St. John, J. B.
High-performance liquid chromatography of metribuzin and non-polar metabolites extracted from leaf tissues 206
Oka, F., see Oka, H. 53
Oka, H.
——, Oka, F. and Ito, Y.
Multilayer coil planet centrifuge for analytical high-speed counter-current chromatography 53
Olsen, L. D.
—— and Hurtubise, R. J.
Mobile phase effects on aromatic hydroxyl compounds with an aminopropyl column and interpretation by the Snyder model 5
Osman, S. F.
—— and Sinden, S. L.
High-performance liquid chromatographic analysis of *Solanum* steroidal alkaloids 189
Oxford, J. M., see Evans, M. B. 170
Packham, A. J., see Fielden, P. R. 117
Palazzolo, D. L.
—— and Quadri, S. K.
Reduced variation in retention times of biogenic amines by temperature control in liquid chromatography with electrochemical detection 216
Parcher, J. F.
—— and Strubinger, J. R.
High-pressure adsorption of carbon dioxide on supercritical-fluid chromatography adsorbents 251
Peltonen, K., see Pukkila, J. 369
Pirkle, W. H.
——, Chang, J.-P. and Burke, III, J. A.
Use of achiral ion-pairing reagents with chiral stationary phases 377
—— and McCune, J. E.
Separation of the enantiomers of N-protected α-amino acids as anilide and 3,5-dimethylanilide derivatives 419
Plicka, J., see Kleinmann, I. 327
Poor, C. L., see Schmitz, H. H. 261
Principi, J. M., see Dubin, P. L. 159
Pukkila, J.
——, Kokotti, H. and Peltonen, K.
High-performance liquid chromatographic determination of benzil in air as an indicator of emissions derived from polyester powder coatings 369
Quadri, S. K., see Palazzolo, D. L. 216
Reed, R. H., see Kerby, N. W. 353
Romero, T. E., see Sánchez-Rasero, F. 424
Rowell, P., see Kerby, N. W. 353
Sánchez-Rasero, F.
——, Romero, T. E. and Dios, C. G.
Determination of ethirimol, in the presence of some normal soil constituents, by liquid chromatography 424
Sandberg, G., see Wingsle, G. 335
Sandra, P., see Janda, V. 200
Sandström, J. F.
——, Skarping, G. and Dalene, M.
Chromatographic determination of amines in biological fluids with special reference to the biological monitoring of isocyanates and amines. II. Determination of 2,4- and 2,6-toluenediamine using glass capillary gas chromatography and selected ion monitoring 135
——, see Skarping, G. 125
Sangö, C., see Skarping, G. 125
Schäfer, K.-H., see Voncken, P. 410
Schepers, G., see Voncken, P. 410
Schmitz, H. H.
——, Artz, W. E., Poor, C. L., Dietz, J. M. and Erdman, Jr., J. W.
High-performance liquid chromatography and capillary supercritical-fluid chromatography separation of vegetable carotenoids and carotenoid isomers 261
Serra, M. A.
——, Avilés, F. X., Giralt, E. and Cuchillo, C. M.
Kinetic analysis of the carboxypeptidase A hydrolysis of oligopeptides by reversed-phase high-performance liquid chromatography 27
Sherman, W. R., see Hsu, F.-F. 437
Siems, W. F., see St. Louis, R. H. 221
Simon, P.
——, Brand, F. and Lemacon, C.
Florisil® sorbent sampling and ion chromatographic determination of airborne aliphatic carboxylic acids 445
Simon, V. A.
—— and Taylor, A.
High-sensitivity high-performance liquid chromatographic analysis of diquat and paraquat with confirmation 153
Sinden, S. L., see Osman, S. F. 189
Sing, K. S. W., see Ahsan, T. 17
Singh, A. K.
——, Gordon, B., Hewetson, D., Granley, K., Ashraf, M., Mishra, U. and Dombrovskis, D.
Screening of steroids in horse urine and plasma by using electron impact and chemical ionization gas chromatography–mass spectrometry 233

Skarping, G.
——, Dalene, M., Brorson, T., Sandström, J. F., Sangö, C. and Tiljander, A.
Chromatographic determination of amines in biological fluids with special reference to the biological monitoring of isocyanates and amines. I. Determination of 1,6-hexamethylenediamine using glass capillary gas chromatography and thermionic specific detection 125
——, see Sandström, J. F. 135
——, see Tiljander, A. 145
Šmídl, P., see Kleinmann, I. 327
Smith, M. S., see Evans, M. B. 170
Steenbeke, G., see Janda, V. 200
St. John, J. B., see Norman, H. A. 206
St. Louis, R. H.
——, Siems, W. F. and Hill, Jr., H. H.
Evaluation of direct axial sample introduction for ion mobility detection after capillary gas chromatography 221
Strubinger, J. R., see Parcher, J. F. 251
Sugitani, H., see Chikazumi, N. 85
Susini, A., see Gülaçar, F. O. 61
Svoboda, V., see Kleinmann, I. 327
Taylor, A., see Simon, V. A. 153
Teale, P., see Houghton, E. 73
Theyson, T. W., see Metz, P. A. 107
Thomas, J.
Analyse par chromatographie liquide haute performance ultra-rapide de la théophylline et de l'anisate de sodium (ou de potassium) dans différentes préparations pharmaceutiques 430
Tiljander, A.
——, Skarping, G. and Dalene, M.
Chromatographic determination of amines in biological fluids with special reference to the biological monitoring of isocyanates and amines. III. Determination of 4,4'-methylenedianiline in hydrolysed human urine using derivatization and capillary gas chromatography with selected ion monitoring 145
——, see Skarping, G. 125
Tressl, R., see Engel, K.-H. 176
Van de Ven, L., see Hetem, M. 269
Van der Leer, T., see Hermans-Lokkerbol, A. 39
Ven, L. van de, see Hetem, M. 269
Verpoorte, R., see Hermans-Lokkerbol, A. 39
Voncken, P.
——, Schepers, G. and Schäfer, K.-H.
Capillary gas chromatographic determination of *trans*-3'-hydroxycotinine simultaneously with nicotine and cotinine in urine and blood samples 410
Walter, H.
—— and Krob, E. J.
Analysis of density-fractionated rat red blood cells of different ages by partitioning in two-polymer aqueous phase systems 307
Warth, L. M.
——, Cooper, R. S. and Fritz, J. S.
Low-capacity quaternary phosphonium resins for anion chromatography 401
Whitman, W. B., see Mesbah, M. 297
Wingsle, G.
——, Sandberg, G. and Hällgren, J.-E.
Determination of glutathione in Scots pine needles by high-performance liquid chromatography as its monobromobimane derivative 335
Yamamoto, E., see Lewis, N. G. 345
Yamazaki, M., see Fujii, Y. 319
Zaikov, Chr., see Bardarov, V. 97

Errata

J. Chromatogr., 469 (1989) 383–389
p. 385, Table I, 13th anion should read "$S_2O_3^{2-}$".
p. 386, 11th line, "Na_2SO" should read "Na_2SO_4".
p. 387, 11th line, reference number 16 should be inserted.